



DR. LEVIN'S POST-OPERATIVE INSTRUCTIONS

Apply **ICE PACKS** (Ziploc bags with peas or gauze soaked in ice water) to your eyelids for 20 minutes every hour while awake for **48 HOURS** after surgery. Icing is **NOT** necessary **AFTER** 48 hours.

Mild oozing is normal in the first 1-2 days. If there is brisk or persistent bleeding, hold firm pressure on the area with a gauze for 10 minutes. If it persists, call the office.

Apply a small amount of the prescribed ointment to the incisions 2 times per day with a Q-tip until the sutures are removed. After the sutures are removed, apply at bedtime for 1 week. If the ointment gets into your eyes, it will make vision blurry.

Sleep with extra 1-2 pillows and keep your head elevated during the day

AVOID dusty or dirty environments and sun exposure (wear large sunglasses). **AVOID** any make-up close to the incisions for 2 weeks. Do **NOT** wear contact lenses for at least 10 days following surgery.

DO NOT touch, rub, or itch your incisions. Take Benadryl if you have itching. If scabs form on the incisions, do not pick at them. Use warm compresses as needed for comfort after the first 48 hours.

You can shower 24 hours after surgery, but avoid water directly on the incisions

AVOID any activities that may increase blood flow to your eyelids or raise your blood pressure for 1 week: bending over, picking up heavy items, sports and sexual relations. Light activities (walking) are fine after day 3.

You will have bruising and swelling. This usually peaks at 48-72 hours. One side may be more swollen and/or bruised. This may extend down to your cheeks due to gravity. The swelling & bruising may last for up to 3-4 weeks (average 1-2 weeks)

Your sutures will be removed 6-8 days after surgery depending on your procedure and how you heal

You may resume driving as soon as you feel that it is safe to do so, usually after 48 hours

CONTACT THE OFFICE IMMEDIATELY IF

- You develop a fever, extreme swelling, or yellow/green discharge from the incision
- There is a sudden loss of vision, double vision or rapidly forming hematoma that is firm and/or results in inability to open the eye
- Bleeding persists despite 10 minutes of firm pressure with gauze
- The incision opens up
- **IF YOU ARE HAVING AN URGENT PROBLEM AND IT IS AFTER HOURS, WEEKEND, OR IF YOU CANNOT REACH ANYONE AT THE OFFICE, PLEASE CALL DR. LEVIN'S CELL (917) 593-4187**

YOUR FOLLOW-UP APPOINTMENT IS ON _____ AT _____